

Monterey B.t. RTU (Ready-To-Use)

- **Biological Insecticide**
- **Controls Worms & Caterpillars on Fruits, Vegetables, Ornamentals & Shade Trees in and around the Home Garden**



Active Ingredient:

Bacillus thuringiensis subspecies *kurstaki* strain SA-12 solids, spores and Lepidopteran active toxins (At least 6 million viable spores per mg)*9.83%
Other Ingredients:90.17%
Total: 100.00%

*The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

EPA Reg. No. 70051-113-54705

EPA Est. No. 70051-CA-001

Net Contents: 1 QT/32 FL OZ/946.36 mL

Distributed by:

LAWN AND GARDEN PRODUCTS, INC.
P.O. Box 35000 • Fresno, CA 93745 • (559) 499-2100
info@montereylawnngarden.com

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Hot Line Number: 1-800-255-3924.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

MONTEREY B.t. RTU:

- Kills worms and caterpillar stage insects, but has no effect on birds, earthworms, or beneficial insects such as honeybees and ladybugs, when used as directed.
- Worms and caterpillars eat treated foliage, then immediately stop feeding and damaging plants.
- Acceptable for use on edible plants up to the day of harvest.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FOR USE

Apply MONTEREY B.t. RTU when worms or caterpillars are first noticed, then repeat at five (5) to seven (7) day intervals while they are active. Apply more frequently to control heavy infestations. Apply thoroughly to top and bottom of foliage. Reapply after heavy rains.

MONTEREY B.t. RTU must be eaten by worms or caterpillars to be effective. After ingesting the insecticide, they immediately stop feeding, though they may otherwise appear to be unaffected for several days. Best results are obtained by treatments when worms are small; they must be actively feeding on treated, exposed foliage. You may apply MONTEREY B.t. RTU up to and on the day of harvest.

HOW TO APPLY

Always shake MONTEREY B.t. RTU thoroughly before use.

Spray leaf surfaces thoroughly, top and bottom, for complete control. Shake regularly while spraying.

FRUITS AND VEGETABLES: Spray all plant foliage. Apply at first sign of infestation and repeat at weekly intervals when needed to insects in the fruit and vegetable garden:

Cabbage Looper, Armyworms, Hornworms, and other Lepidopteran (moth) pests on broccoli, cauliflower, collards, kale, mustard greens, turnip greens, cabbage, celery, lettuce, melons, and tomatoes.

SHADE TREES & ORNAMENTALS in and around the Home: Spray at first sign of infestation and repeat at weekly intervals when needed to maintain control. Apply to thoroughly cover all foliage surfaces of shade trees and ornamentals for control of the following leaf-feeding worms:

Bagworm	Spring Cankerworm
Fall Cankerworm	Gypsy Moth*
Tent Caterpillar	Elm Spanworm
Fall Webworm	

*Apply when buds swell, but before leaves are fully open. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first appearance.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool place. Activity may be impaired by storage at temperatures above 90°F.

PESTICIDE DISPOSAL and CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

For containers less than or equal to 5 gallons:

If empty: Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

For containers greater than 5 gallons:

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

WARRANTY

Lawn and Garden Products, Inc. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the Directions for Use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OF MERCHANTABILITY IS MADE.

(CER-082812A / 110413N) 0513/1113(02)