

DR. IRON[®] RTU HOUSEPLANT FOOD

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Information concerning the raw materials composing this product can be obtained by writing to Lawn and Garden Products, Inc. and referring to the product name.

NOTICE: Because weather, crop, soil, and other conditions may vary, Lawn and Garden Products, Inc. and/or the Seller make no warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risks of use or handling, whether or not in accordance with directions.

0410(01)

- **Ready-To-Use**
- **Promotes Rapid Greenup and Dark Green Color**
- **Apply Indoors & Outdoors**
- **Low Stain Formula**



GUARANTEED ANALYSIS **0.16 – 0 – 0**

| | |
|--|-------|
| Total Nitrogen (N) | 0.16% |
| 0.16% Water Soluble Organic Nitrogen (N) | |
| Sulfur (S) | 0.04% |
| Iron (Fe) | 0.06% |

Derived from Lo-Biuret Urea and Ferrous Sulfate.

DR. IRON is a registered trademark of Lawn and Garden Products, Inc.

ATTENTION: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

NET CONTENTS: 1 Quart / 946 mL

Guaranteed By:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745 • (559) 599-2100
www.montereylawngarden.com

**KEEP OUT OF REACH OF CHILDREN
CAUTION: HARMFUL IF SWALLOWED.**

GENERAL INFORMATION

DR. IRON[®] RTU HOUSEPLANT FOOD is designed to promote rapid green-up and dark green color. DR. IRON[®] RTU HOUSEPLANT FOOD is most effective when plants are growing rapidly and least effective when plants are dormant or suffering from stress.

PRECAUTIONS

- Shake container well before using.
- Store above 50°F.
- Do not apply foliar spray to plants stressed from moisture or heat.
- If spilled on concrete, wash off immediately.

DIRECTIONS FOR USE

Shake bottle thoroughly prior to using. Apply to houseplants or individual plants. Spray to thoroughly wet. Do not over apply. Repeat application on a 2 week schedule when plants are actively growing. Avoid spraying directly onto flower blossoms or buds.