

NOTE: This is a specimen label for electronic distribution. Always refer to product label on container for specific directions for use.



FUNGI-FIGHTER

SYSTEMIC FUNGICIDE

*For Control of Certain Diseases
of Flowers, Foliage Plants, Shrubs, Turf, Grapes,
Apples, Melons, and Shade Trees.
Prevents and Cures Diseases.*

FOR HOMEOWNER USE ONLY

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-(H-1,2,4-triazol-1-yl)-2-butanone 25%

INERT INGREDIENTS: 75%

Total 100%

EPA Reg. No. 54705-3

EPA Est. No. 11160-CA-1

Manufactured For:
LAWN AND GARDEN PRODUCTS, INC.
P. O. Box 35000 • Fresno, CA 93745
(559) 499-2100

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If In Eyes: Flush with plenty of water. Call a physician if irritation persists.

If On Skin: Wash with plenty of soap and water. Get medical attention.

SYMPTOMS OF POISONING: The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by hyperactivity followed by sedation.

NOTE TO PHYSICIANS: No specific antidote. Treat symptomatically.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Causes eye irritation. Prolonged or frequently repeated skin contact may cause allergic skin reaction in some individuals. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Do not make applications when weather conditions favor drift from target area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in original container in cool, dry place out of reach of children and domestic animals. Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection.

NOTICE

Seller makes no warranty, express or implied, concerning use other than indicated on label. Buyer assumes all responsibility for safety and use not in accordance with directions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For homeowner use only. Read entire label and use strictly in accordance with label precautionary statements and directions.

APPLICATION: FUNGI-FIGHTER is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage is necessary. Rainfall or sprinkler irrigation, within 1/2 hour after application, does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Applications should be applied at recommended intervals to maintain disease control.

TURF DISEASE CONTROL		
Disease	Rate* Oz./1,000 sq. ft. in 2-4 gallons of water	Remarks
Anthracnose Brown Patch/Rhizoctonia Blight Copper Spot Dollar Spot Powdery Mildew Red Thread Rusts	1 - 2	Preventative: Apply at 15-30 day intervals prior to appearance of the disease. Curative: Apply the higher rate when disease is apparent. Follow up with the preventative rate.
Southern Blight	1 - 4	
Stripe Smut	2	Apply a single application in the spring as grass growth begins.
Fusarium Blight Summer Patch	2 - 4	Apply in the spring 30-60 days before symptoms appear. Repeat at 30-day intervals.
Gray Snow Mold /Typhula Blight (except California) Pink Snow Mold/ Fusarium Patch (except California)	2 - 4	Make one application in late fall.

ORNAMENTAL PLANT DISEASE CONTROL			
Plant	Disease	Rate*	Remarks
Azalea Dendrobium (Hawaii only)	Flower blight	1 - 2 teaspoons/ 5 gallons of water	Apply to the point of drip. Repeat at 7 - 21 day intervals.
Cedar Iris Nephthytis Zinnia	Leaf blight		
Azalea Begonia Crape Myrtle Euonymous Geranium Hollyhock Oak Rose Snapdragon Sweetpea Sycamore Viburnum Willow Zinnia	Powdery mildew		
Geranium Hollyhock Pine Snapdragon	Rusts		
Pine	Tip Blight		

GREENHOUSE PLANTS (Homeowner only)			
Plant	Disease	Rate*	Remarks
African Violet Carnation Chrysanthemum Geranium Grapeleaf Ivy Rose	Powdery mildew	1 - 2 teaspoons/ 5 gallons of water	Apply to the point of drip. Repeat at 30-day intervals.
Carnation Chrysanthemum Geranium	Rust		

* 1 oz. = 3-1/3 tablespoons of FUNGI-FIGHTER
3 level teaspoons = 1 level tablespoon of FUNGI-FIGHTER

NOTE: This is a specimen label for electronic distribution. Always refer to product label on container for specific directions for use.

HOME GROWN FRUIT, MELONS and VEGETABLE DISEASE CONTROL			
Plant	Disease	Rate*	Remarks
Apple Pear	Cedar-apple rust (apple only) Powdery mildew	½ - 2 teaspoons/ 5 gallons of water	Apply in adequate water to the point of drip. Make the first application to apples at green-tip stage and to pears at the budburst stage; continue at 7-14 day intervals. Do not exceed 6 teaspoons/season. May be applied up to 45 days before harvest.
Grape	Powdery mildew	1-3 teaspoons/ 5 gallons of water	Apply pre-bloom and continue at 14-21 day intervals.
	Black rot	1-1½ teaspoons/ 5 gallons of water 1½-2 teaspoons/ 5 gallons of water	Preventative: Apply first application at 10 inch green shoot and continue at 7-14 day intervals until berry coloring is complete. Post-infection: During bloom, apply within 72 hours of the infection period. After bloom, continue on a 7-14 day schedule. Do not exceed 9 teaspoons/season. May be applied up to 14 days before harvest.
Cucurbits Balsam Pear Chinese Waxgourd Citrus Melon Cucumber Gourd Pumpkin Squash Watermelon	Powdery mildew	2-4 teaspoons/ 5 gallons of water	Apply to the point of drip. Begin application when disease first appears and repeat at 14-day intervals. Do not exceed 16 teaspoons/season. May be used up to 1 day before harvest.

* 1 oz. = 3-1/3 tablespoons of FUNGI-FIGHTER
3 level teaspoons = 1 level tablespoon of FUNGI-FIGHTER