

REMUDA FULL STRENGTH

41% Glyphosate Concentrate With Surfactant

Complete Broad Spectrum Post Emergence Herbicide
For Vegetation Management, Ornamental Weed Control
and Non-Planted Areas

Controls Many Annual and Perennial Grasses and Broadleaf Weeds



Active Ingredient:

Glyphosate, N-(phosphonomethyl) glycine, in
the form of its isopropylamine salt* 41.0%

Inert Ingredients: 59.0%
Total 100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

EPA Reg. No. 228-366-54705

EPA Est. No. 48498-CA-1

NET CONTENTS: _____

Distributed By:

LAWN AND GARDEN PRODUCTS, INC.

P.O. Box 35000 • Fresno, CA 93745 • (559) 499-2100

www.montereylawnandgarden.com

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed or inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist. Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

USER SAFETY RECOMMENDATIONS

Users Should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. DO NOT mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Not for use on (1) turf being grown for sale or other commercial use as sod; (2) trees being grown for sale or other commercial use, or for the production of timber or wood products; (3) turf, trees, crops, timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants in non-crop and non-timber only.

GENERAL INFORMATION

REMUDA FULL STRENGTH, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most hand held standard sprayers after dilution and thorough mixing with water in accordance with label instructions.

REMUDA FULL STRENGTH moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

DOMESTIC ANIMALS: REMUDA FULL STRENGTH is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

NOTE: This is a specimen label for electronic distribution. Always refer to product label on container for specific directions for use.

ATTENTION

AVOID DRIFT. Extreme care must be used when applying this product to prevent injury to desirable plants.

DO NOT allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the plant or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals, or other unintended consequences. Keep container closed to prevent spills and contamination.

APPLICATION INSTRUCTIONS

Hand Held Equipment: Use Coarse Sprays Only.

Mix REMUDA FULL STRENGTH in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed on this label, apply a 0.5% solution of REMUDA FULL STRENGTH to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. Allow three or more days before tillage or mowing.

For annual weeds over 6 inches tall, use a 1% solution. For best results, use a 2% solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods which result in less than complete coverage, use a 5% solution for annual and perennial weeds and a 5-10% solution for woody brush and trees.

Prepare the desired volume of spray solution by mixing the amount of REMUDA FULL STRENGTH in water as shown in the following table:

Desired Volume	Amount of REMUDA FULL STRENGTH				
	0.5%	1%	2%	5%	10%
1 Gallon	2/3 Oz.	1-1/3 Oz.	2-2/3 Oz.	6-1/2 Oz.	13 Oz.

2 Tablespoons = 1 Fluid Ounce

WEEDS CONTROLLED

REMUDA FULL STRENGTH controls many annual and perennial grasses and broadleaf weeds, such as:

ANNUAL WEEDS

- | | |
|------------------|-------------------------|
| Balsamapple | Panicum |
| Bassia, fivehook | Ragweed, common |
| Brome | Ragweed, giant |
| Fiddleneck | Smartweed, Pennsylvania |
| Fleabane, hairy | Sowthistle, annual |
| Fleabane | Sunflower |
| Kochia | Thistle, Russian |
| Lettuce, prickly | Velvetleaf |

Apply to actively growing grass and broadleaf weeds. Allow at least 3 days after treatment before tillage. For maximum benefit, apply when weeds are 6 inches or less in height. To prevent seed production, applications should be made prior to seedhead formation. REMUDA FULL STRENGTH does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

Extreme care must be exercised to avoid contact of spray with foliage, green stems, exposed non-woody roots, desirable turfgrasses, trees, shrubs or other desirable vegetation since severe damage or destruction may result.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds. Do not exceed 7.8 fluid ounces per 1,000 square feet per year.

PERENNIAL WEEDS

- | | | |
|----------------------|--------------------------|---------------------|
| Alfalfa | Dock, curly | Pampasgrass |
| Alligatorweed* | Dogbane, hemp | Paragrass |
| Anise (fennel) | Fescues | Phragmites* |
| Artichoke, Jerusalem | Fescue, tall | Poison hemlock |
| Bahiagrass | Guineagrass | Quackgrass |
| Bentgrass | Horsenettle | Redvine* |
| Bermudagrass | Horseradish | Reed, giant |
| Bindweed, field | Ice plant | Ryegrass, perennial |
| Bluegrass, Kentucky | Johnsongrass | Smartweed, swamp |
| Blueweed, Texas | Kikuyugrass | Spurge, leafy* |
| Brackenfern | Knapweed | Starthistle, yellow |
| Bromegrass, smooth | Lantana | Sweet potato, wild* |
| Bursage, woollyleaf | Lespedeza | Thistle, artichoke |
| Canarygrass, reed | Milkweed | Thistle, Canada |
| Cattail | Muhly, wirestem | Timothy |
| Clover, red | Mullein, common | Torpedograss* |
| Clover, white | Napiegrass | Trumpet creeper* |
| Cogongrass | Nightshade, silverleaf | Vaseygrass |
| Dallisgrass | Nutsedge; purple, yellow | Velvetgrass |
| Dandelion | Orchardgrass | Wheatgrass, western |

* Partial control

Apply REMUDA FULL STRENGTH as follows to control or destroy most perennial weeds. NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages. Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Extreme care must be exercised to avoid contact of spray with foliage, green stems, exposed non-woody roots, desirable turfgrasses, trees, shrubs or other desirable vegetation since severe damage or destruction may result.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds. Do not exceed 7.8 fluid ounces per 1,000 square feet per year.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: Store above 10°F (-12°C) to keep product from crystallizing. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and shake, roll or agitate to mix well before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

CONTAINER DISPOSAL: Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer and the limit of liability of any such use. The exclusive remedy of user or buyer and the limit of liability of Lawn and Garden Products, Inc, is the purchase price paid for the quantity of product involved.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

(RV 022503B)
0901/0404(03)