

STRAWBERRY & FRUIT TREE WEEDER

Preemergence Herbicide for Weed Control



ACTIVE INGREDIENT: Napropamide 2.0%
 INERT INGREDIENTS: 98.0%
 TOTAL: 100.0%

Contains 0.02 Pounds of Active Ingredient per 1 Pound.

EPA Reg. No. 70506-33-54705

EPA Est. No. 48498-CA-1

NET CONTENTS: 7 Lbs.

Manufactured For:
LAWN AND GARDEN PRODUCTS, INC.
 P. O. Box 35000 • Fresno, CA 93745 • (559) 499-2100
 www.montereylawnandgarden.com

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes, and clothing. May be harmful if swallowed. Do not breathe dust. Avoid contamination of food and feed.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves made of any waterproof material such as Barrier Laminate, Nitrile Rubber (> 14 mils), Neoprene rubber (> 14 mils) or Viton (> 14 mils).
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

GENERAL PRECAUTIONS

- Read all instructions before using.
- Use only for recommended purposes and at recommended rates.
- Do not overdose.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product as specified on this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated areas until dusts have settled.

WHAT THIS PRODUCT DOES

STRAWBERRY & FRUIT TREE WEEDER controls certain annual grassy and broadleaf weeds as they germinate. It does not control established weeds. The weeds controlled are listed below:

WEEDS CONTROLLED BY STRAWBERRY & FRUIT TREE WEEDER

Annual Grasses:

- Annual Bluegrass
- Wild Barley
- Barnyardgrass, Junglerice
- Bromes
 - Cheatgrass
 - Downy Brome
 - Ripgut Brome
 - Soft Chess
- Large (Hairy) Crabgrass
- Smooth Crabgrass
- Southwestern Cupgrass
- Foxtails
 - Bristly Foxtail
 - Giant Foxtail
 - Green Foxtail
 - Yellow Foxtail

- Goosegrass
- Seedling Johnsongrass
- Wild Oats
- Panicums
 - Guineagrass
 - Fall Panicum
 - Texas Panicum
 - Witchgrass
- Italian Ryegrass
- Sandbur
- Red Sprangletop
- Stinkgrass

Annual Broadleaf Weeds:

- Carpetweed
- Common Chickweed
- Coast Fiddleneck
- Filaree
- Groundsel
- Prostrate Knotweed
- Lambsquarters
- Prickly Lettuce

- Little Mallow*
- Redroot Pigweed
- Pineappleweed
- Common Purslane
- Horse Purslane
- Common Ragweed**
- Annual Sowthistle

- * From seed
- ** Suppression

WHERE TO USE THIS PRODUCT

STRAWBERRY & FRUIT TREE WEEDER can be applied over small ornamental plants or around larger plants in rose gardens, container plantings, ornamental landscapes and ground covers including dichondra.

NOTE: This is a specimen label for electronic distribution. Always refer to product label on container for specific directions for use.

Use STRAWBERRY & FRUIT TREE WEEDER for weed control in the following plants:

Fruits and Nuts

Almond	Lemon	Pistachio
Apple	Nectarine	Plum
Apricot	Orange	Prune
Cherry	Peach	Tangerine
Filbert	Pear	Walnut
Grape	Pecan	

Strawberry: Applications should be delayed until the desired number of daughter plants have become established. DO NOT apply from bloom through harvest.

Deciduous and Evergreen Trees

Acer	Eucalyptus	Picea
Ash	Ficus	Pin Oak
Birch	Fir	Pinus
Black Walnut	Gleditsia	Pittosporum
Cedar	Hemlock	Podocarpus
Citrus	Japanese Larch	Populus
Crataegus	Laburnum	Prunus
Cypress	Malus	Pears
Dogwood	Olive (<i>Olea</i>)	Pistachios
Douglas Fir	Palm (<i>Washingtonia robusta</i>)	

Deciduous and Evergreen Shrubs

Abelia	Forsythia	Oleander
Asparagus (<i>Sprengeri</i>)	Gardenia	Osmanthus
Aucuba	Heather (<i>Calluna</i>)	Photinia
Azalea	Hibiscus	Pittosporum
Bottlebrush	Honeysuckle (<i>Lonicera</i>)	Podocarpus
Bougainvillea	Hypericum	Pyracantha
Buxus	Ilex	Raphiolepis
Camellia	Juniperus	Rhododendron
Cotoneaster	Leucothe	Rosa
Crepe Myrtle	Ligustrum	Star Jasmine
Dogwood	Nandina	Taxus
Euonymus	Natal Plum	Viburnum
		Xylosema

Groundcovers

Ajuga	Hedera	Liriop
Carpobrotus	Hypericum	Osteosperma
Delosperma	Iceplant	Pachysandra
Dichondra	Ivy	Sedum
Erysimum	Lantana	Vinca
Gazania		

Flowers

African Daisy	Dahlia	Narcissus
Aster	Daisy	Petunia
Bird of Paradise	Geranium	Plantain Lily (<i>Hosta</i>)
Chrysanthemum	Gladiolus	Zinnia

WHEN TO USE THIS PRODUCT

STRAWBERRY & FRUIT TREE WEEDER should be used before weeds germinate. It will not control established weeds.

Application can be made anytime of the year to weed-free soil. Fall and winter applications will provide excellent weed control the following spring and summer.

In containers, apply after setting and watering the plants.

STRAWBERRY & FRUIT TREE WEEDER should be spread evenly on the soil surface using a shaker can or granular herbicide applicator.

DO NOT APPLY when weather conditions favor drift from areas treated.

When weed infestation is heavy or when broadleaves are the primary weeds, use 6.9 lbs./1,000 sq. ft. For annual grassy weeds or when weed infestation is light, use 4.6 lbs./1,000 sq. ft.

DO NOT APPLY more than 4.6 lbs. STRAWBERRY & FRUIT TREE WEEDER per 1,000 sq. ft. in California.

STRAWBERRY & FRUIT TREE WEEDER should be incorporated evenly into the soil using a hand cultivator to a depth of 1-2 inches or irrigation or natural moisture (rainfall or snow) (1/2 inch within 2-3 days) for optimum results.

For weed control in dichondra, apply at the time of seeding or to establish dichondra including dichondra sod. Irrigate with a minimum of one inch of water.

RECOMMENDATIONS: TURFGRASS

(For Use on Warm Season Grasses Only)

STRAWBERRY & FRUIT TREE WEEDER is recommended for annual bluegrass, crabgrass and goosegrass control in the following established turfgrasses: bahia, bermudagrass, centipede, St. Augustine and fescue.

TIMING OF APPLICATION:

For control of crabgrass and goosegrass, apply STRAWBERRY & FRUIT TREE WEEDER at the recommended rate. Application should be made 2-3 weeks prior to weed emergence. STRAWBERRY & FRUIT TREE WEEDER should be applied with a gravity flow drop spreader or a cyclone type spreader. A more uniform application can be made by spreading half the required amount over the area to be treated, then applying the remainder at right angles.

STRAWBERRY & FRUIT TREE WEEDER requires a minimum of one half (1/2) inch of rainfall or irrigation 2-3 days after application for adequate control. If irrigation methods are not available, the timing of the application of STRAWBERRY & FRUIT TREE WEEDER should be moved back several weeks to allow for some rainfall between application and weed germination.

RATES OF APPLICATION:

For Crabgrass Control: Apply 2.25 - 3.38* lbs. of STRAWBERRY & FRUIT TREE WEEDER uniformly per 1,000 sq. ft. Use the higher rate when applying before January 1 (not applicable north of South Carolina or when weed infestation is expected to be heavy).

*Where a rate range is given, use the higher rate when weed infestation is expected to be heavy.

For Goosegrass Control: Apply 2.25 - 3.38 lbs. of STRAWBERRY & FRUIT TREE WEEDER per 1,000 sq. ft. Goosegrass usually starts to germinate 6-7 weeks after the start of crabgrass germination.

Split Application for Crabgrass and Goosegrass Control: When both of these weeds are to be controlled, apply 2.25 lbs. of STRAWBERRY & FRUIT TREE WEEDER spread uniformly per 1,000 sq. ft. for crabgrass and another 2.25 lbs. of STRAWBERRY & FRUIT TREE WEEDER per 1,000 sq. ft. for goosegrass. Apply 8-10 weeks apart. Timing should be adjusted to the germination of weeds in your area. Split applications give longer and better weed control than one application.

For Annual Bluegrass (*Poa Annua*) Control: Annual bluegrass is a winter annual in most of the warm season turf regions; therefore, application of STRAWBERRY & FRUIT TREE WEEDER should be made in the fall before the annual bluegrass germinates. Apply 2.25 - 3.38 lbs. STRAWBERRY & FRUIT TREE WEEDER per 1,000 sq. ft.

Reseeding: DO NOT reseed or overseed within six months after application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container out of the reach of children and pets. In case of leak or spill, contain material and dispose of as waste.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Lawn and Garden Products, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Lawn and Garden Products, Inc. harmless for any claims relating to such factors.

Lawn and Garden Products, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Lawn and Garden Products, Inc., and Buyer and User assume the risk of any such use. LAWN AND GARDEN PRODUCTS, INC. MAKES NO IMPLIED WARRANTY OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

In no event shall Lawn and Garden Products, Inc. or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LAWN AND GARDEN PRODUCTS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LAWN AND GARDEN PRODUCTS, INC., THE REPLACEMENT OF THE PRODUCT.

Lawn and Garden Products, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Lawn and Garden Products, Inc.

1203(01)