

# Monterey Consan® 20

## Disinfectant • Deodorizer • Mildewstat

- Cleans, Disinfects and Deodorizes on Hard, Non-porous Surfaces
- Eliminates Odors Caused By Bacteria, Mold and Mildew
- Helps Inhibit the Growth of Unsightly Algae (Not for use in CA.)
- Controls Listed Fungal Diseases in Turf
- Controls Fairy Rings and Toad Stools in Turf (Not for use in CA.)

### ACTIVE INGREDIENTS:

Alkyl (60% C14, 30% C16, 5% C12, 5% C18) Dimethyl Benzyl Ammonium Chloride . . . . . 10.0%  
Alkyl (68% C12, 32% C14) Dimethyl Ethylbenzyl Ammonium Chloride . . . . . 10.0%

**OTHER INGREDIENTS:** . . . . . 80.0%

**TOTAL:** . . . . . 100.0%

EPA Reg. No.: 10324-94-54705

EPA Est. No.: 48498-CA-1

See inside booklet for Complete Precautionary Statements, First Aid, Directions for Use & Storage and Disposal

### KEEP OUT OF REACH OF CHILDREN

## DANGER PELIGRO

SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE.  
IF YOU DO NOT UNDERSTAND THE LABEL, FIND SOMEONE TO EXPLAIN IT TO YOU IN DETAIL.

### FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

**IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.

Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER.** Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and chemical-resistant gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear coveralls worn over long-sleeved shirt and long pants, waterproof gloves, chemical-resistant footwear and socks, protective eyewear, chemical-resistant headgear when using this product for algae control in overhead watering system and chemical-resistant apron when mixing, loading or cleaning equipment. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates.

**Net Contents:**

1 Pint  1 Quart  1 Gallon

### DISTRIBUTED BY:

**Lawn and Garden Products, Inc.**

PO Box 35000 • Fresno, CA 93745 • (559) 499-2100 • info@montereylawnngarden.com

Made in the USA

Consan is a registered trademark of Brandt Consolidated, Inc.

**GENERAL DISINFECTION PERFORMANCE:** This product kills the following bacteria in 10 minutes at 1.5 oz. per 5 gal. of water (469 ppm active) and 5% soil on hard, non-porous surfaces: *Campylobacter jejuni*, *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Proteus mirabilis*, *Salmonella enterica*, *Staphylococcus aureus*, *Staphylococcus aureus* Methicillin Resistant (MRSA), *Yersinia enterocolitica*.

**MILDEWSTATIC PERFORMANCE:** This product controls the following mold at 2 oz. per 5 gal. of water (625 ppm active) and 5% soil on hard, non-porous surfaces: *Aspergillus niger*.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

Rinse hard, non-porous food contact surfaces with potable water after application of product.

**AGRICULTURAL USE REQUIREMENTS:** Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this section (box) only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls worn over long sleeved shirt and long pants, waterproof gloves, chemical-resistant shoes plus socks, and protective eyewear.

**NON-AGRICULTURAL USE REQUIREMENTS:** The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets out of the treated area until sprays have dried.

### FOR USE AS A GENERAL DISINFECTANT ON HARD, NON-POROUS SURFACES:

1. Pre-clean heavily soiled areas.
2. Apply 1.5 oz. of this product per 5 gal. of water (or equivalent use dilution) to hard, non-porous surfaces using a sponge, brush, cloth, mop, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray.
3. Treated surfaces must remain wet for 10 minutes.
4. Wipe dry or allow to air dry.
5. Prepare a fresh solution daily or when visibly dirty.

**TO CONTROL MOLD/MILDEW:** Pre-clean hard, non-porous surfaces. Prepare use solution by adding 2 oz. of this product per 5 gal. of water (or equivalent use dilution). Apply use solution to hard, non-porous surfaces which will effectively inhibit the growth of mold and mildew and their odors. Repeat treatment every seven days, or more often if new growth appears.

**GENERAL DEODORIZATION:** To deodorize, apply 1.5 oz. of this product per 5 gal. of water to hard, non-porous surfaces. Wipe up excess liquid and allow to air dry.

### WORK AREAS AND BENCHES, POTS, FLATS AND FLOWER BUCKETS, CUTTING TOOLS:

Pre-clean surfaces. Spray or swab hard, non-porous working surfaces or soak cutting edge of tool with a solution of 1.5 oz. of this product per 5 gallons of water (or equivalent use dilution) before each work period and again after each plant is completed to help control transfer of diseases such as *Botrytis*, crown rot, downy mildew, *Erwinia* and root rot<sup>†</sup>. Allow surface to remain wet for 10 minutes. To apply solution as a sprayer application, use as a coarse spray only and spray 6-8 inches from surface. Do not breathe spray. Wipe up excess liquid or allow to air dry. Prepare a fresh solution daily or when visibly dirty. (†Not for use in CA.)

### TREATMENT OF TURF:

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. (Use against dollar spot (*Sclerotinia*), fusarium blight (*Fusarium*), Southern blight (*Sclerotium rolfsii*), rust, toad stools and fairy rings not allowed in CA.) To control fungal diseases including dollar spot (*Sclerotinia*), fusarium blight (*Fusarium*), leaf spot (*Helminthosporium*), brown patch (*Rhizoctonia*), fading out (*Curvularia*), Southern blight (*Sclerotium rolfsii*), pythium blight, rust, toad stools and fairy rings and algal build-up in turf grasses, including lawns of grass or Dichondra, spray a solution of 0.5 oz. of this product per gal. of water to treat 50 square feet (5 ft. x 10 ft.) of lawn. Treat entire lawn to prevent spread of infection at least one foot beyond the edges of isolated diseased areas. Saturate the entire treatment area, wetting all plant and soil surfaces. Repeat spray applications every 10 - 14 days, as needed to control new or established disease conditions. Treatment is required during the warm growing season and must be made when the lawn is dry or damp, but not during wet rainy weather.

**GREENHOUSE GLASS** (Not for use in CA.): Spray or swab hard, non-porous surfaces with a solution of 1.5 oz. of this product per 5 gallons of water (or equivalent use dilution). Allow to air dry. Wash off dead algae with water. Spray clean surface again with the solution. Do not rinse. For sprayer applications, use as a coarse spray only. Allow surfaces to remain wet for 10 minutes then air dry. Prepare a fresh solution daily or when visibly dirty.

**HOUSEPLANT AND GARDENING CONTAINERS AND EQUIPMENT** (Not for use in CA.): To control algae and cross contamination in terrariums, bubbles, and other hard, non-porous gardening surfaces, wipe surface with a solution of 1.5 oz. of this product per 5 gallons of water (or equivalent use dilution). Allow surfaces to remain wet for 10 minutes then air dry. Prepare a fresh solution daily or when visibly dirty.

**WALKWAYS** (Not for use in CA.): This product will inhibit the growth of algae on hard, non-porous walkways. For heavy infestations, spray or swab surface with a solution of 1.5 oz. of this product per 5 gallons of water (or equivalent use dilution). Let stand for an hour or more then brush and wash away dead algae. Soak area again with the solution. Do not rinse. Allow to dry on the surface and repeat application when algae growth returns. For sprayer applications, use as a coarse spray only. Allow surfaces to remain wet for 10 minutes then air dry. Prepare a fresh solution daily or when visibly dirty.

**FOR ALGAE TREATMENT OF CONTAINERIZED FOUNTAINS, WATER DISPLAYS, DECORATIVE POOLS/PONDS AND STANDING WATER:** DO NOT use when fish or other wildlife are present. *This product is not to be used in open water ways connected to larger watersheds or in waters that serve as natural habitats for aquatic or amphibious organisms. Spray from fountains treated with this product will not harm nearby plantings.*

Remove floating type algae by raking, dragging with cable or chain, skimming, or any other suitable method. It is more economical to remove floating type algae before the water is treated.

**Initial Dosage:** Add 5 oz. of this product per 1,562 gal. of water (or equivalent use dilution) at any point that is convenient, such as the bowl, pool, or sump.

**Weekly Maintenance:** Add 2.5 oz. of this product per 1,562 gal. of water (or equivalent use dilution) every 3 – 5 days or as to maintain 0.5 ppm active. Each month drain and clean bowl. Refill with fresh water and repeat initial treatment. Draining removes airborne dirt, dust, contamination and alkali build-up.

**Note:** Algae does not usually grow in water that is 5 feet or more in depth. However, floating type algae may grow on pond weeds that grow on the surface of the water.

**FOUNTAIN AND POOL TREATMENT DILUTION TABLE:**

Fountain or Pool Capacity gal. of Water	Initial Treatment	Maintenance Dose Booster Dose
195	0.6 oz.	0.3 oz.
390	1.2 oz.	0.6 oz.
781	2.5 oz.	1.3 oz.
1,562	5.0 oz.	2.5 oz.

**SANITIZATION OF EXTERIOR HOUSEHOLD SURFACES DIRECTIONS**

**PREPARATION OF USE SOLUTION:** Apply a use solution of 0.13 oz. of this product per gal. of water (or equivalent use dilution) to sanitize hard, non-porous exterior surfaces such as vinyl, plastic, sealed concrete, painted or sealed woodwork, and sealed stucco. Surfaces to be treated include house siding, decks, patios, walkways and driveways. One-half gal. of diluted product will treat 200-300 sq. ft. of hard, non-porous surfaces.

**APPLICATION:** Pre-clean heavily soiled areas. Apply solution with a brush, mop, cloth, sponge, hand pump trigger spray device, coarse trigger spray device, so as to wet surfaces thoroughly. For sprayer applications, spray 6 inches to 8 inches from surface. Do not breathe spray. If using a pressure washer with high-pressure spray to sanitize hard, non-porous surfaces, wear suitable respiratory protective equipment and protective eyewear to control exposure to spray. Treated surfaces much remain wet for 3 minutes.

**WATER AND SMOKE DAMAGE RESTORATION**

**SEWER BACKUP & RIVER FLOODING** (Not for use in CA.): To use as a deodorizer, dilute 0.5 to 1 oz. of this product per gal. of water allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affected areas with a mechanical spray device, coarse pump or trigger spray device before and after cleaning and extraction. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Use proper ventilation; open windows.

**CARPETS, CARPET CUSHIONS, UPHOLSTERY, DRAPES AND OTHER POROUS MATERIALS, SUB FLOORS, DRYWALL, TRIM AND FRAME LUMBER, TACKLESS STRIP AND PANELING** (Not for use in CA.): To use as a deodorizer against water damage, extract the excess water. Test hidden area for colorfastness. Dilute 0.5 to 1 oz. of this product per gal. of water, allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil. Apply directly with a mechanical spray device or coarse pump or trigger spray device to fully saturate affected materials. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Roll, brush or agitate into materials. Follow with a thorough extraction. Dry rapidly and thoroughly.

**WATER AND SMOKE DAMAGE RESTORATION** (Not for use in CA.): Effective against odor caused by smoke and water damage for home, institutional, industrial and hospital use. This product is particularly suitable for use in water damage restoration. Dilute 0.5 to 1 oz. of this product per gal. of water, allowing for the diluting effect of absorbed water within saturated materials. Saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

**COMMERCIAL ORCHID, HORTICULTURAL AND ORNAMENTAL USES**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State and Tribe, consult the State/Tribal agency responsible for pesticide regulation.

**EVAPORATIVE COOLER** (Not for use in CA.): Controls algae, slime forming fungi on cooler pads and certain fungal plant pathogens that may carry over in the cooler as well as bacterial odors. Use 1 teaspoon of this product per 15 gallons of water every other week.

**Note:** Not all varieties or strains of plants listed have been tested. Unintentional consequences such as crop injury or ineffectiveness may result because of certain environmental or growing conditions, manner of use or applications, and insect resistance. Therefore, before treating a large number of plants, spray a few plants and observe for plant damage and for performance prior to full-scale application. All risks shall be assumed by the user.

## ORCHIDS

Species	Diseases Controlled	Rate/Gal of water	Additional Information
Seedlings	Damping-off (caused by Pythium, Fusarium, or Rhizopus)	0.2 oz.	From flask, soak seedling in solution for 2-5 minutes before potting to remove agar. Spray seedlings once a month as preventative maintenance. Spray any time disease is evident with solution.
Dividing plants		0.5 oz.	Remove compost then soak plant for 2-5 minutes in solution. Operate with wet hands and cutting tools (see Disinfecting). After division and root trim, soak divisions in solution for 2 to 5 minutes. Pot in compost drenched with solution.
Adult Plants		0.25 oz.	Spray any time disease, other than rot (see Adult Orchid Rots), is evident with solution. Spray once per month as preventative maintenance. Drench compost when fungus is evident.
	Botrytis petal blight	0.02 oz.	During cool weather, spray plants, flowers and under benches at first sign of spotting on flowers. Normally, weekly sprayings will greatly reduce or eliminate the botrytis spotting.
	Rots (such as <i>Erwinia</i> , et al)	0.5 oz.	Remove compost from plant, then soak entire plant 10 minutes or more in solution. Cut away all rot with treated tool (see Disinfecting). Soak plant wounds again for 10 minutes in solution. Repot in drenched compost.
	Crown rot	0.25 oz.	Spray plants weekly with solution, assuring coverage around and in base of leaves.

## HORTICULTURAL USAGES

Species	Diseases Controlled	Rate/Gal of water	Additional Information
Bulbs such as Daffodils, Easter lilies, Gladioli, Hyacinth, Iris, Narcissus, Tulips	Fusarium basal rot	0.5 oz.	Pre-storage: Treat the corms and bulbs immediately after they are "cleaned" by soaking them for 10 minutes in the solution.  Pre-plant: Treat again immediately before planting.
Cut Flowers in containers, tubes and decorative vases	Bacterial stem plugging (restricts water uptake) Fungi (can create ethylene gas, a case of petal drop)	0.13 oz.	Flower spikes should be hardened in solution during harvesting, storage and shipping to help control the transfer of diseases.
Plant Cuttings such as Begonias, Carnations, Chrysanthemums, Fuchsias, Geraniums)	Carnation rust, Crown gall, Gray mold leaf spots and Powdery mildew	0.25 oz.	Soak cuttings 2-5 minutes in the solution. Plant in compost drenched with the solution.
Seedlings	Damping off	0.25 oz.	Treat seeds and cuttings before planting by spraying with solution to completely wet all surfaces. Use inert material for planting seeds and cuttings, such as sphagnum moss, vermiculite, perlite, pumice, sterilized commercial mix rather than garden soil. If garden soil must be used, thoroughly saturate with the solution.  After planting, water with solution (unless using soil saturated as above) and again after seedlings emerge. Spray weekly as needed.
	Downymildew	0.25 oz.	Spray with solution at 5 day intervals as needed.

## ORNAMENTAL USAGES

Species	Diseases Controlled	Rate/Gal of water	Additional Information
Aucuba, Dahlia, Fuchsia, Geranium, Gladiolus, Peony, Piersis	Botrytis	0.25 oz.	Spray buds and blooms lightly at first sign of disease, and repeat 2-3 times per week as needed. Before shipping, dip spikes in solution.
	Downy mildew, Gray mold	0.25 oz.	Spray at first sign of disease repeating at 7-14 day intervals. Before shipping, dip spikes in solution.
	Rust	0.25 oz.	Spray at 7-14 day intervals during the wet periods starting in early Summer. Before shipping, dip spikes in solution.
Begonias, Ferns, Gesneria	Bud rot, Leaf spots, Gray mold, Shoot rot	0.13 oz.	Spray solution before buds open, when blossoms show pink, at ¼ blossom fall, and 2 weeks after petal fall.
Cactus	Bacterial leaf spot, Root rot, Top rot	0.25 oz.	Sterilize seeds and disease free cuttings by dipping cuttings in the solution for 5 minutes before planting. Then spray at 10-14 day intervals.
	Slimy collar rot	0.25 oz.	Just before planting, dip entire plant in the solution.
Calendulas, China Asters, Chrysanthemum, Dahlias, Lilacs, etc.	Powdery mildew	0.25 oz.	Spray upper and lower surface of leaves and branches with solution until thoroughly wet. Spray when buds burst in Spring and repeat in 2 weeks. Spray every 2 weeks during the mildew season.
Carnations	Canker rot, Carnation rust, Crown rot, Leaf spot, Root rots, Stem rot	0.25 oz.	Spray at first sign of disease and repeat at 7-10 day intervals.
	Gray mold rot	0.25 oz.	Spray 1-2 times just before bloom and again as flowers are opening.
Crassula, Kalanchoe, Sedum	Crown gall, Crown rot, Root rot	0.25 oz.	Destroy infected plant parts and drench the remainder. Spray plants weekly as needed.
Dogwood	Anthrachnose, Leaf blight, Leaf spots	0.25 oz.	Spray 4 times: when flower buds begin to open, as soon as flower bracts have fallen, 4 weeks later, and in late summer after flower buds are well formed. If possible, spray just before rainy period.
Fuchsia, Gardenia, Geranium, Heliotrope	Bud rot, Gray mold, Stem rot	0.25 oz.	Spray at first sign of disease and repeat weekly as needed.
Hawthorn	Rust	0.25 oz.	Begin protective spraying with new leaf formation after winter dormancy but before onset of rainy season. Thoroughly spray undersides of infected leaves.
Hawthorn (Crataegus)	Leaf blights	0.25 oz.	Make 3 applications: begin when leaves emerge from buds, when leaves are half grown, and again when fully grown.
Hollyhock	Rust	0.25 oz.	Apply at weekly intervals as soon as growth begins. Repeat applications in late Summer.
Iris	Botrytis, Crown rot, Gray mold rot	0.25 oz.	Treat all new rhizomes before planting. Submerge rhizomes and 3-4" of leaves in the solution for 30 minutes. Plant at once while still wet with solution.
	Leaf spots	0.25 oz.	Spray weekly with solution while leaves are growing.
Ornamental trees (such as Sycamore, Ash, Crape Myrtle, Live Oak, etc.)	Anthrachnose	0.25 oz.	Spray upper and under sides of leaves and twigs until thoroughly wet. A large tree requires 50-60 gallons of spray per treatment for adequate coverage. Make 3 applications: when buds start to swell in the spring, as leaves start to expand, and when leaves are about 3/4 grown.
	Powdery Mildew	0.25 oz.	Spray upper and lower surface of leaves and branches with solution until thoroughly wet at bud break and again in April. Apply 50-60 gallons of spray per tree for adequate coverage.

ORNAMENTAL USAGES (Continued)			
Species	Diseases Controlled	Rate/Gal of water	Additional Information
Palms	Heart rot, Penicillium, leaf base rot	0.25 oz.	At first sign of rot, completely fill cup or "heart" formed by the palm branches at top center of the plant. Repeat every 7 days as needed.
Roses	Black spots, Powdery mildew, Rusts, Stem gall	0.25 oz.	When symptoms appear, spray upper and lower surface of leaves and branches with solution until thoroughly wet. Apply at 7-10 day intervals as needed.
	Brown canker	0.25 oz.	Begin spray applications in early spring when new growth starts and repeat weekly until plant is dormant.
Rubber Plant	Anthrachnose, Crown gall	0.13 oz.	Spray plant at first sign of disease and repeat weekly as needed.
Saintpaulias (African Violets)	Bud rot, Crown rot, Downy mildew, Gray mold rot (Botrytis), Powdery mildew, Root rot	0.09 oz.	Dip entire plant and pot in the solution for 5 minutes. Thereafter, spray every 10 days as needed.
Snapdragons	Anthrachnose, Downy mildew, <i>Phyllosticta</i> blight, Powdery mildew, Stem rot, Wilt	0.25 oz.	When symptoms appear, spray upper and lower sides of leaves until thoroughly wet. Repeat every 7-10 days as needed. Collect and burn tops in the fall.
	Rust	0.25 oz.	Apply at 7-10 day intervals as soon as plants are set out. Collect and burn tops in the Fall.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

### CONDITIONS OF SALE AND WARRANTY

Lawn and Garden Products, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in accordance with the use directions under normal conditions. To the extent consistent with applicable law, Lawn and Garden Products, Inc. neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability. The directions for use of this product are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of Lawn and Garden Products, Inc. and Seller. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, Lawn and Garden Products, Inc. or the Seller shall not be liable for consequential, special, indirect or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Lawn and Garden Products, Inc. and is accepted as such by the Buyer.

(MSN-092117A) A-18 1115/1017 (02)

BATCH NO.